

## Activity study of possible endocrine disruption via the TRb or PPARy2 receptors by using CALUX cell systems

**Commonly used acronym:** TRb and PPARy2 CALUX

Created on: 03-05-2022 - Last modified on: 05-05-2022

### Organisation

**Name of the organisation** Sciensano

**Department** Chemical and physical health risks

**Country** Belgium

### Partners and collaborations

BDS

## SCOPE OF THE METHOD

<b>The Method relates to</b>	Human health
<b>The Method is situated in</b>	Basic Research
<b>Type of method</b>	In vitro - Ex vivo
<b>Specify the type of cells/tissues/organs</b>	U2-OS cells (human osteoblasts), stably transfected with the receptor and a luciferase reporter construct

## DESCRIPTION

### Method keywords

cell culture  
CALUX  
reporter gene assays

### Scientific area keywords

Endocrine disruptors  
obesogens  
PPARy  
TRb  
mixtures  
concentration addition

### Method description

The CALUX® system (Chemically Activated LUCiferase eXpression) of Bio Detection Systems (BDS, Amsterdam, the Netherlands) uses U-2 OS cells (human osteoblast) that are stably transfected with human TRb or human PPARg2 (BDS, Amsterdam) and a luciferase reporter construct under the control of a receptor specific response element.

Through measuring the activity of chemical compounds on these cell lines, we can determine if they can potentially have endocrine activity. The activity of mixture of chemicals can also be determined in these cell systems.

### Lab equipment

Cell incubator,  
Safety cabinet,  
Fluorimeter.

### Method status

Published in peer reviewed journal

## PROS, CONS & FUTURE POTENTIAL

### Advantages

Speed (vs *in vivo* experiments),  
Useful for prioritisation.

### Challenges

Difficult to extrapolate to the *in vivo* situation because of Absorption, Distribution, Metabolism, and Excretion (ADME).

## REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

### References

Mertens, B, Van Heyst A, Demaegdt H, Boonen I, Van Den Houwe K, Gosciny S, Elskens M, and Van Hoeck E. Assessment of hazards and risks associated with dietary exposure to mineral oil for the Belgian population. Food and Chemical Toxicology, 2021, Volume 149

I Boonen I, Van Heyst A , Van Langenhove K, Van Hoeck E, Mertens M, Denison MS, Elskens M, Demaegdt H. Assessing the receptor-mediated activity of PAHs using AhR-, ER?- and PPAR?- CALUX bioassays. Food Chem Toxicol. 2020, 145:111602. doi: 10.1016/j.fct.2020.111602

Simon C, Onghena M, Covaci A, Van Hoeck E, Van Loco J, Vandermarken T, Van Langenhove K, Demaegdt H, Mertens B, Vandermeiren K, Scippo ML, Elskens M. Screening of endocrine activity of compounds migrating from plastic baby bottles using a multi-receptor panel of *in vitro* bioassays.

Mertens B, Van Hoeck E, Blaude MN, Simon C, Onghena M, Vandermarken T, Van Langenhove K, Demaegdt H, Vandermeiren K, Covaci A, Scippo ML, Elskens M, Van Loco J. Evaluation of the potential health risks of substances migrating from polycarbonate replacement baby bottles. Food Chem Toxicol. 2016, 97:108-119.

Demaegdt H, Daminet B, Evrard A, Scippo ML, Muller M, Pussemier L, Callebaut A, Vandermeiren K. Endocrine activity of mycotoxins and mycotoxin mixtures. Food Chem Toxicol. 2016, 96:107-16.

Pereira-Fernandes A, Demaegdt H, Vandermeiren K, Hectors TL, Jorens PG, Blust R, Vanparys C. Evaluation of a Screening System for Obesogenic Compounds: Screening of Endocrine Disrupting Compounds and Evaluation of the PPAR Dependency of the Effect. PLoS One. 2013, 8 : e77481.

Coordinated by



Financed by



Vlaanderen  
verbeelding werkt

