

Identifying Key Genes involved in Drug-Induced Liver Injury using Machine Learning on Human in vitro data sets

Created on: 14-02-2024 - Last modified on: 15-02-2024

Organisation

Name of the organisation Vrije Universiteit Brussel (VUB)

Department In Vitro Toxicology and Dermato-Cosmetology

Specific Research Group or Service In Vitro Toxicology and Dermato-Cosmetology

Country Belgium

Geographical Area Brussels Region

SCOPE OF THE METHOD

The Method relates to	Human health
The Method is situated in	Translational - Applied Research
Type of method	In silico

DESCRIPTION

Method keywords

Feature selection

Machine learning

Supervised classification
Wrapper feature selection
transcriptomic signatures
machine learning algorithms
prediction models
Toxicogenomics
artificial intelligence

Scientific area keywords

Drug-induced cholestasis
drug development
preclinical drug toxicity testing
Hepatotoxicity
hepatic toxicity

Method description

Drug-induced intrahepatic cholestasis (DIC) is a main type of hepatic toxicity that is challenging to predict in early drug development stages. Preclinical animal studies often fail to detect DIC in humans. *In vitro* toxicogenomics assays using human liver cells have become a practical approach to predict human-relevant DIC. The present study was set up to identify transcriptomic signatures of DIC by applying machine learning algorithms to the Open TG-GATEs database. A total of nine DIC compounds and nine non-DIC compounds were selected, and supervised classification algorithms were applied to develop prediction models using differentially expressed features. Feature selection techniques identified 13 genes that achieved optimal prediction performance using logistic regression combined with a sequential backward selection method. The internal validation of the best-performing model showed accuracy of 0.958, sensitivity of 0.941, specificity of 0.978, and F1-score of 0.956. Applying the model to an external validation set resulted in an average prediction accuracy of 0.71. The identified genes were mechanistically linked to the adverse outcome pathway network of DIC, providing insights into cellular and molecular processes during response to chemical toxicity. Our findings provide valuable insights into toxicological responses and enhance the predictive accuracy of

DIC prediction, thereby advancing the application of transcriptome profiling in designing new approach methodologies for hazard identification.

Lab equipment

- Open TG-GATEs database,
- GEO database,
- Affymetrix GeneChip,
- Classification algorithms.

Method status

Internally validated

Published in peer reviewed journal

PROS, CONS & FUTURE POTENTIAL

Advantages

- Such a signature could aid in *in vitro* DIC prediction, facilitating early detection of this chemical-induced toxicity.
- The identified features have biologically interpretable functions, mechanistically anchored in an AOP network, and provide new insights into molecular and cellular behavior processes during DIC development, making them valuable tools for understanding and predicting toxicological responses.

Challenges

Differential expression (DE) analysis often generates numerous correlated candidate genes, leading to redundant information and reduced translatability for laboratory testing and lowered translatability of the DE findings for high-throughput laboratory testing. To address this, a permutation-based approach was employed to refine the results that estimate feature relevance by measuring changes in model performance upon permuting feature vectors .

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

Links

[Unraveling the mechanisms underlying drug-induced cholestatic liver injury: ide...](#)

Other remarks

This work was performed in the context of the ONTOX project (<https://ontox-project.eu/>) supported by the European Commission (grant agreement 963845) and as part of the ASPIS project cluster (<https://aspis-cluster.eu/>).

Coordinated by



Financed by



Vlaanderen
verbeelding werkt

