

LipidTOX assay in primary rat hepatocytes

Commonly used acronym: LipidTOX

Created on: 05-03-2019 - Last modified on: 28-02-2022

Contact person

Kaat Leroy

Organisation

Name of the organisation Vrije Universiteit Brussel (VUB)

Department Pharmaceutical and Pharmacological Sciences

Specific Research Group or Service In Vitro Toxicology and Dermato-Cosmetology

Country Belgium

Geographical Area Brussels Region

SCOPE OF THE METHOD

The Method relates to	Human health
The Method is situated in	Basic Research, Translational - Applied Research
Type of method	In vitro - Ex vivo
Species from which cells/tissues/organs are derived	Rat
Type of cells/tissues/organs	Primary rat hepatocytes

DESCRIPTION

Method keywords

Steatosis
Phospholipidosis
Hepatocytes
neutral lipids
phospholipids
Hepatotoxicity

Scientific area keywords

Hepatotoxicity
cytotoxicity
Steatosis
Phospholipidosis

Method description

The method detects two facets of drug-induced cytotoxicity i.e. the intracellular accumulation of phospholipids and of neutral lipids, i.e. phospholipidosis and steatosis respectively. The assay makes use of a kit containing an aqueous, red-fluorescent formulation of labelled phospholipids (LipidTOX™ Red phospholipid stain, excitation/emission ~595/615 nm) which is up taken by the cells upon incubation with a phospholipidosis-inducing compound. The second component of the kit is a selective green-fluorescent stain for neutral lipids (LipidTOX™ Green neutral lipid stain, excitation/emission ~495/505 nm), which can be used sequentially on fixed cells for the analysis of steatosis or can be used independently for single-parameter analysis (Nioi et al. 2007). Additionally, use of VECTASHIELD® Mounting Medium containing 4',6-diamidino-2-phenylindole (DAPI) which binds directly to DNA and produces upon excitation a blue fluorescence, enables intracellular localisation of the lipids.

Lab equipment

Fluorescence microscope

Method status

History of use

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

References

- Anderson N. and Borlak J. (2006) "Drug-induced phospholipidosis." *FEBS Letters* 580: 5533-40
- Elaut G., Henkens T., Papeleu P., Snykers S., Vinken M., Vanhaecke T. and V. Rogiers (2006) "Molecular mechanisms underlying the dedifferentiation process of isolated hepatocytes and their cultures." *Current Drug Metabolism* 7: 629-60
- Gomez-Lechon M.J., Lahoz A., Gombau L., Castell J.V. and Donato M.T. (2010) "In vitro evaluation of potential hepatotoxicity induced by drugs." *Current Pharmaceutical Design* 16: 1963-77
- Kodavanti U.P. and Mehendale H.M. (1990) "Cationic amphiphilic drugs and phospholipid storage disorder." *Pharmacological Reviews* 42:327-54
- McKim J.M., (2010) "Building a tiered approach to in vitro predictive toxicity screening: a focus on assays with in vivo relevance." *Combinatorial Chemistry and High Throughput Screening* 13: 188-206
- Monteith D.K., Morgan R.E. and Halstead B. (2006) "In vitro assays and biomarkers for drug-induced phospholipidosis." *Expert Opinion on Drug Metabolism and Toxicology* 2: 687-96
- Morgan R.E., Kriauciunas A., Berridge B., Sullivan J. and Monteith D.K. (2004) "An in vitro surrogate for drug-induced phospholipidosis." *Toxicologist* 78: 388
- Nioi P., Perry B.K., Wang E.J, Gu Y.Z. and R.D. Snyder (2007) "In vitro detection of drug-induced phospholipidosis using gene expression and fluorescent phospholipid based methodologies." *Toxicological Sciences* 99: 162-73
- Papeleu P., Vanhaecke T., Henkens T., Elaut G., Vinken M., Snykers S. and Rogiers V. (2006) "Isolation of rat hepatocytes." *Methods in Molecular Biology* 320: 229-37
- Reasor M.J., Hastings K.L. and Ulrich R.G. (2006) "Drug-induced phospholipidosis: issues and future directions." *Expert Opinion on Drug Safety* 5:567-83
- Sawada H., Takami K. and Asahi S. (2005) "A toxicogenomic approach to drug-induced phospholipidosis: analysis of its induction mechanism and establishment of a novel in vitro screening system." *Toxicological Sciences* 83:282-92

Sawada H., Taniguchi K. and Takami K. (2006) "Improved toxicogenomic screening for drug-induced phospholipidosis using a multiplexed quantitative gene expression ArrayPlate assay." *Toxicology In Vitro* 20:1506-13

Seglen, P.O. (1976) "Preparation of isolated rat liver cells." *Methods in Cell Biology* 13:29-83

Coordinated by



Financed by



Vlaanderen
verbeelding werkt



bruxelles
environnement
leefmilieu
brussel
.brussels