

# Cytotoxicity measurement in cultured primary rat hepatocytes

*Commonly used acronym: MTT assay*

*Created on: 11-03-2019 - Last modified on: 28-02-2022*

## Contact person

Axelle Cooreman

## Organisation

**Name of the organisation** Vrije Universiteit Brussel (VUB)

**Department** Pharmaceutical and Pharmacological Sciences

**Specific Research Group or Service** In Vitro Toxicology and Dermato-Cosmetology

**Country** Belgium

**Geographical Area** Brussels Region

## SCOPE OF THE METHOD

<b>The Method relates to</b>	Human health
<b>The Method is situated in</b>	Basic Research
<b>Type of method</b>	In vitro - Ex vivo
<b>Species from which cells/tissues/organs are derived</b>	rat
<b>Type of cells/tissues/organs</b>	primary hepatocytes

## DESCRIPTION

### Method keywords

cell viability test

MTT cytotoxicity assay  
cytotoxicity of chemicals  
mitochondria  
Succinate dehydrogenase  
Formazan

### **Scientific area keywords**

Cell culture  
cell viability  
Primary hepatocytes  
rat  
liver

### **Method description**

The MTT test is performed to determine the *in vitro* cytotoxicity of selected chemicals. The mitochondrial enzyme succinate dehydrogenase is responsible for the biotransformation of toxic agents and MTT. The ability of cells to reduce MTT provides an indication of the mitochondrial integrity and activity, which in turn may be interpreted as a measure of viability and/or cell number. When chemical compounds are induced in primary rat hepatocytes, their cell viability and their possibility to transform xenobiotical substances decreases. In this respect, when the MTT solution (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) is added to the exposed cells, the possibility to transform this pale yellow salt into dark blue formazan crystals decreases. The formazan crystals formed in the cells are solubilized in DMSO and can be measured colorimetrically.

### **Lab equipment**

Multiplate reader.

### **Method status**

History of use

## **PROS, CONS & FUTURE POTENTIAL**

### **Advantages**

Easy to apply;  
Not so many extra materials or solutions needed;  
The method itself is very fast (4h).

## Challenges

The density of the cells should be even in each well. Make sure that the cell suspension is homogenous and devoid large cell aggregates.

There are possible interferences between the tested chemical and the MTT as substrate. MTT can be directly reduced by test substances and give artifacts.

Therefore, before initiating experiments, a special procedure that allows quantification of the "true" MTT mitochondrial reduction from the "false" chemical MTT reduction should be performed.

Considerable cell death is observed shortly after isolation of hepatocytes from a freshly removed liver and during the early phases of cultivation. To reduce the effects of this experimentally-induced cell injury and thus to avoid false positive results, experiments should be initiated at earliest 24h after cells seeding.

## REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

### References

Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods* 65: 55-63

Xu G., Yan Z., Wang N. and Liu Z. (2011) Synthesis and cytotoxicity of cis-dichloroplatinum (II) complexes of (1S,3S)-1,2,3,4-tetrahydroisoquinolines. *European Journal Medicinal Chemistry* 46(1): 356-363

Porto I.C., Oliveira D.C., Raelle R.A., Ribas K.H., Montes M.A. and De Castro C.M. (2011) Cytotoxicity of current adhesive systems: In vitro testing on cell cultures of primary murine macrophages. *Dental Materials* 27(3): 221-228

Vinken M., Decrock E., De Vuyst E., Leybaert L., Vanhaecke T. and Rogiers V. (2009) Biochemical characterisation of an in vitro model of hepatocellular apoptotic cell death. *Alternatives to Laboratory Animals* 37: 209-218

Coordinated by



Financed by

