

Flow-cytometric determination of neutral lipids

Created on: 20-03-2019 - Last modified on: 28-02-2022

Contact person

Joost Boeckmans

Organisation

Name of the organisation Vrije Universiteit Brussel (VUB)

Department Pharmaceutical and Pharmacological Sciences

Specific Research Group or Service In Vitro Toxicology and Dermato-Cosmetology

Country Belgium

Geographical Area Brussels Region

SCOPE OF THE METHOD

The Method relates to	Human health
The Method is situated in	Translational - Applied Research
Type of method	In vitro - Ex vivo
Specify the type of cells/tissues/organs	Human skin-derived adults stem cells

DESCRIPTION

Method keywords

Flow-cytometry neutral lipids quantitative fluorimetric in vitro

Scientific area keywords

Steatosis stem cells lipids lipid accumulation

Method description

Using this method you can measure the (relative) lipid load in human skin-derived stem cells differentiated towards hepatic cells. This method could also be applied on other cell types (e.g. HepG2), since it is based on the following publication: "M. T. Donato et al., Chem. Biol. Interact. 181, 417–423 (2009)." Briefly: 1. Aspirate medium from the cell

culture 2. Incubate 10' with TrypLE (200 μ L/well for 24- multiwell format) 3. Add 500 μ L pre-warmed PBS (37°C) to every well and harvest the sample 4. Rinse with 500 μ L PBS 5. Centrifigate according to the cell type 6. Resuspend in 1 mL PBS (containing BODIPY(TM) 1:2500 (see publication above)) on ice 7. 5' before measuring + 1 μ L Hoechst (+homogenize by pipetting) 8. Dilute 1:10 (PBS (4 °C)) before measuring to limit background signal 9. Measure signal (up to 100.000 events)

Lab equipment

Flow-cytometer; Cell culture equipment; Biosafety cabinet.

Method status

Internally validated

PROS, CONS & FUTURE POTENTIAL

Advantages

Fast.

Challenges

Measuring many samples can be time-consuming.

Modifications

You can use also other cell types.

Future & Other applications

Drug testing (anti-steatotic drugs); Assessing drug-induced liver steatosis (*in vitro*).

REFERENCES, ASSOCIATED DOCUMENTS AND OTHER INFORMATION

References

M. T. Donato et al., Chem. Biol. Interact. 181, 417–423 (2009)

R. M. Rodrigues et al., Stem Cells Dev. 23, 44–55 (2014)

Coordinated by







